

# Effect of Door-to-Balloon Time on Mortality in Patients With ST-Segment Elevation Myocardial Infarction

Robert L. McNamara, MD, MHS,\* Yongfei Wang, MS,\* Jeph Herrin, PhD,\* Jephtha P. Curtis, MD,\* Elizabeth H. Bradley, PhD,† David J. Magid, MD, MPH,§|| Eric D. Peterson, MD, MPH,¶|| Martha Blaney, PHARM D,# Paul D. Frederick, PhD,\*\* Harlan M. Krumholz, MD, SM,\*††††† for the NRMI Investigators

*New Haven, Connecticut; Denver, Colorado; Durham, North Carolina; South San Francisco, California; and Seattle, Washington*

**OBJECTIVES** We sought to determine the effect of door-to-balloon time on mortality for patients with ST-segment elevation myocardial infarction (STEMI) undergoing primary percutaneous coronary intervention (PCI).

**BACKGROUND** Studies have found conflicting results regarding this relationship.

**METHODS** We conducted a cohort study of 29,222 STEMI patients treated with PCI within 6 h of presentation at 395 hospitals that participated in the National Registry of Myocardial Infarction (NRMI)-3 and -4 from 1999 to 2002. We used hierarchical models to evaluate the effect of door-to-balloon time on in-hospital mortality adjusted for patient characteristics in the entire cohort and in different subgroups of patients based on symptom onset-to-door time and baseline risk status.

**RESULTS** Longer door-to-balloon time was associated with increased in-hospital mortality (mortality rate of 3.0%, 4.2%, 5.7%, and 7.4% for door-to-balloon times of  $\leq 90$  min, 91 to 120 min, 121 to 150 min, and  $>150$  min, respectively;  $p$  for trend  $<0.01$ ). Adjusted for patient characteristics, patients with door-to-balloon time  $>90$  min had increased mortality (odds ratio 1.42; 95% confidence interval [CI] 1.24 to 1.62) compared with those who had door-to-balloon time  $\leq 90$  min. In subgroup analyses, increasing mortality with increasing door-to-balloon time was seen regardless of symptom onset-to-door time ( $\leq 1$  h,  $>1$  to 2 h,  $>2$  h) and regardless of the presence or absence of high-risk factors.

**CONCLUSIONS** Time to primary PCI is strongly associated with mortality risk and is important regardless of time from symptom onset to presentation and regardless of baseline risk of mortality. Efforts to shorten door-to-balloon time should apply to all patients. (J Am Coll Cardiol 2006;47: 2180–6) © 2006 by the American College of Cardiology Foundation

Time to reperfusion for patients with ST-segment elevation myocardial infarction (STEMI) consistently predicts mortality for fibrinolytic therapy (1–3). In contrast, studies have found conflicting results regarding the relationship between mortality and time to reperfusion with primary percutaneous coronary intervention (PCI). Some investigators have found lower mortality for shorter symptom onset-to-reperfusion time for all patients (4) or just certain subgroups such as high-risk patients (5) or those presenting within 2 h of symptom onset (6). Other studies found no lower mortality for shorter symptom onset-

to-balloon time but did find lower mortality for shorter door-to-balloon time (7,8). Finally, some studies failed to find an association between either symptom onset-to-balloon time or door-to-balloon time and mortality (9,10).

Although the American College of Cardiology/American Heart Association (ACC/AHA) guidelines for management of patients with STEMI recommend door-to-balloon times of 90 min or less (11,12), a minority of patients are currently treated within this time period, and this pattern has not changed recently (13). The perception that time to reperfusion is less important in PCI (9,10) may contribute to the current inertia in performance. To evaluate the effect of door-to-balloon time on mortality in these patient groups, we used detailed patient-level and hospital-level longitudinal data from a national sample of patients with STEMI admitted from 1999 to 2002 from the National Registry of Myocardial Infarction (NRMI)-3 and -4 (14).

## METHODS

**Study design and sample.** We used NRMI, a voluntary acute myocardial infarction (AMI) registry sponsored by

From the \*Department of Medicine, Section of Cardiovascular Medicine, †Department of Epidemiology and Public Health, Section of Health Policy and Administration, and ‡Robert Wood Johnson Clinical Scholars Program, Yale University School of Medicine, New Haven, Connecticut; §Clinical Research Unit, Kaiser Permanente, Denver, Colorado; ||Departments of Emergency Medicine and Preventive Medicine and Biometrics, University of Colorado Health Sciences Center, Denver, Colorado; ¶Duke Clinical Research Institute, Duke University, Durham, North Carolina; #Genentech Inc., South San Francisco, California; \*\*Ovation Research Group, Seattle, Washington; and the ††Center for Outcomes Research and Evaluation, Yale-New Haven Hospital, New Haven, Connecticut. This research was supported by the National Heart, Lung, and Blood Institute, R01HL072575. Dr. Blaney is employed by Genentech, Inc.

Manuscript received August 12, 2005; revised manuscript received December 15, 2005, accepted December 19, 2005.

#### Abbreviations and Acronyms

ACC	= American College of Cardiology
AHA	= American Heart Association
AMI	= acute myocardial infarction
CI	= confidence interval
ECG	= electrocardiogram
NRMI	= National Registry of Myocardial Infarction
PCI	= percutaneous coronary intervention
STEMI	= ST-segment elevation myocardial infarction

Genentech Inc. (South San Francisco, California), to define a cohort of patients with STEMI who received acute reperfusion therapy with primary PCI. The NRMI criteria (15,16) include a diagnosis of AMI according to the *International Classification of Diseases, Ninth Revision, Clinical Modification* (code 410.X1) and any one of the following criteria: total creatine kinase or creatine kinase-MB that was two or more times the upper limit of the normal range or elevations in alternative cardiac markers; electrocardiographic evidence of AMI or nuclear medicine testing, echocardiography, or autopsy evidence of AMI. During our study period of January 1, 1999, to December 31, 2002, there were 830,473 AMI admissions in NRMI. The following patients were excluded sequentially: patients transferred to or from another acute care institution (n = 341,730); with neither ST-segment elevation (2+ leads) nor left bundle branch block on the first electrocardiogram (ECG) (n = 334,013); with AMI symptom onset after the admission date and time (n = 4,305); with a nondiagnostic first ECG (e.g., the first ECG did not show ST-segment elevation or left bundle branch block; n = 14,314); with diagnostic ECG that preceded hospital presentation by more than 1 h (prehospital ECG), or with time from door-to-diagnostic ECG that was more than 6 h or missing (n = 6,467); who did not receive primary PCI (n = 92,772); with door-to-balloon times that were negative, more than 6 h, or missing (n = 925); and with unknown time of symptom onset (n = 4,804). In addition, to avoid including hospitals that performed primary PCI uncommonly, patients treated in hospitals reporting fewer than 20 PCI patients over the four-year time period (n = 1,921) were excluded. The final cohort included 29,222 patients from 395 hospitals. Mortality status at the time of discharge was known for all patients.

**Data collection and measures.** Our outcome was in-hospital mortality, and the principal independent variable was door-to-balloon time, which is the time from hospital arrival to balloon inflation, derived from the corresponding date/time noted in the medical record and recorded in the NRMI case report form. Patients were stratified based on their time from symptom onset-to-door time ( $\leq 1$  h,  $>1$  to 2 h,  $>2$  h) and whether they had ACC/AHA high-risk factors (anterior/septal location, diabetes mellitus, heart rate  $>100$  beats/min, systolic blood pressure  $<100$  mm Hg) (11).

Other patient-level variables included age ( $<65$  years, 65 to 79 years,  $\geq 80$  years), gender, race/ethnicity (white, black, Hispanic, other), insurance status, and clinical characteristics. Clinical characteristics consisted of medical history (current smoker, chronic renal insufficiency, previous AMI, hypertension, family history of coronary artery disease, hypercholesterolemia, congestive heart failure, previous percutaneous transluminal coronary angioplasty, previous coronary artery bypass graft surgery, chronic obstructive pulmonary disease, stroke, angina, diabetes); presentation characteristics (time from symptom onset-to-presentation, whether a prehospital ECG was performed, the admission time of day [day, evening, or night], admission day of week [weekday or weekend], chest pain at presentation, systolic blood pressure, heart rate, heart failure); and the results of the diagnostic ECG (number of leads with ST-segment elevation, AMI location, ST-segment depression, nonspecific ST/T-wave changes, Q-wave). Calendar time, measured as the number of days between January 1, 1999, and the hospital admission date, was included as an independent variable to account for any secular trends as well as for differing reporting periods by hospitals.

**Statistical analysis.** We first examined the bivariate association between patient characteristics and in-hospital mortality, using chi-square tests to assess for the association between categorical variables and in-hospital mortality and *t* tests or F tests to assess for the association between continuous variables and in-hospital mortality.

We then examined the bivariate association between door-to-balloon time and in-hospital mortality with door-to-balloon time as a categorical variable. We did this for the whole cohort and stratified by symptom onset-to-door time ( $\leq 1$  h,  $>1$  to 2 h,  $>2$  h) and presence or absence of anterior/septal location, diabetes mellitus, heart rate  $>100$  beats/min, systolic blood pressure  $<100$  mm Hg, and any of these baseline risk factors.

For the independent effect of door-to-balloon time on in-hospital mortality, we used a multivariable logistic regression model using in-hospital death as the dependent variable. Because NRMI enrolls hospitals that then report patients, we could not assume that measurements were independent of hospital; assessment of intraclass correlations indicated that variation in both time to treatment ( $p = 0.1099$ , 95% CI 0.0916 to 0.1282) and mortality ( $p = 0.0084$ , 95% CI 0.0052 to 0.0434) was partly explained by hospital. Thus, we used hierarchical models to account for clustering of patients within hospitals. Random effects were specified for the main intercept and the coefficients of calendar time in the model. We replicated the model in all the strata of symptom onset-to-door time and baseline risk factors, as defined above; the stratification variable was not included in the corresponding subgroup model. We also estimated a final set of models using the whole cohort, each of which included the interaction between door-to-balloon time and one of these stratification variables. We performed secondary analyses that included the 2.0% of patients

**Table 1.** Patient Characteristics, Door-to-Balloon Time, and In-Hospital Mortality

Description	Prevalence n (%)	Door-to-Balloon Time		Mortality	
		Median, min (Quartile Range)	p	n (%)	p
All	29,222 (100.00)	102 (54)		1,329 (4.55)	
Demographics					
Age			<0.0001		<0.0001
<65 yrs	17,314 (59.25)	99 (52)		356 (2.06)	
65-80 yrs	9,425 (32.25)	106 (56)		623 (6.61)	
>80 yrs	2,483 (8.50)	115 (59)		350 (14.10)	
Gender					
Male	20,712 (70.88)	100 (52)	<0.0001	739 (3.57)	<0.0001
Female	8,510 (29.12)	108 (57)		590 (6.93)	
Race			<0.0001		0.7847
White	25,082 (85.83)	101 (53)		1,147 (4.57)	
Black	1,428 (4.89)	118 (63)		58 (4.06)	
Hispanic	971 (3.32)	114 (61)		47 (4.84)	
Other	1,741 (5.96)	102 (52)		77 (4.42)	
Health insurance			<0.0001		<0.0001
Medicare only	5,218 (17.86)	107 (58)		397 (7.61)	
Medicare and (commercial or other)	5,037 (17.24)	107 (57)		381 (7.56)	
Medicare and Medicaid	473 (1.62)	113 (62)		51 (10.78)	
Commercial	13,483 (46.14)	100 (52)		313 (2.32)	
Medicaid only	751 (2.57)	104 (56)		37 (4.93)	
Veterans Administration	156 (0.53)	95 (55)		6 (3.85)	
Other	1,302 (4.46)	98 (53)		49 (3.76)	
Self	2,502 (8.56)	97 (50)		80 (3.20)	
Unknown	300 (1.03)	105 (61)		15 (5.00)	
Medical history					
Current smoker	11,320 (38.74)	99 (52)	<0.0001	282 (2.49)	<0.0001
Chronic renal insufficiency	698 (2.39)	117 (65)	<0.0001	109 (15.62)	<0.0001
Diabetes mellitus	5,440 (18.62)	110 (59)	<0.0001	380 (6.99)	<0.0001
Previous myocardial infarction	4,800 (16.43)	106 (59)	<0.0001	237 (4.94)	0.1565
Hypertension	14,218 (48.66)	105 (55)	<0.0001	749 (5.27)	<0.0001
Hypercholesterolemia	11,065 (37.87)	101 (54)	0.0124	327 (2.96)	<0.0001
Family history of coronary artery disease	8,490 (29.05)	101 (53)	0.0005	230 (2.71)	<0.0001
Congestive heart failure	913 (3.12)	118 (65)	<0.0001	126 (13.80)	<0.0001
Percutaneous coronary intervention	4,259 (14.57)	104 (57)	0.0032	150 (3.52)	0.0005
Coronary artery bypass	1,698 (5.81)	120 (65)	<0.0001	109 (6.42)	0.0001
Chronic obstructive pulmonary disease	2,319 (7.94)	107 (58)	<0.0001	152 (6.55)	<0.0001
Stroke	1,357 (4.64)	114 (65)	<0.0001	149 (10.98)	<0.0001
Angina	2,312 (7.91)	105 (58)	0.0288	123 (5.32)	0.0633
Presentation					
Prehospital 12-lead ECG	2,915 (9.98)	83 (50)	<0.0001	103 (3.53)	0.0056
Chest pain at presentation					<0.0001
No	1,527 (5.23)	123 (73)	<0.0001	336 (22.00)	
Yes	27,375 (93.68)	101 (52)	<0.0001	937 (3.42)	
Unknown	320 (1.10)	111 (61)	<0.0001	56 (17.50)	
Systolic blood pressure			<0.0001		<0.0001
<100 mm Hg	2,963 (10.14)	96 (51)		452 (15.25)	
100-180 mm Hg	23,433 (80.19)	102 (54)		786 (3.35)	
>180 mm Hg	2,724 (9.32)	112 (57)		53 (1.95)	
Unknown	102 (0.35)	117 (69)		38 (37.25)	
Heart rate			<0.0001		<0.0001
<50 beats/min	1,766 (6.04)	95 (48)		143 (8.10)	
50-100 beats/min	23,969 (82.02)	102 (53)		793 (3.31)	
>100 beats/min	3,376 (11.55)	113 (64)		355 (10.52)	
Unknown	111 (0.38)	118 (65)		38 (34.23)	
First assessment of heart failure			<0.0001		<0.0001
No congestive heart failure	25,908 (88.66)	101 (53)		722 (2.79)	
Rales/jugular venous distension	1,895 (6.48)	110 (60)		195 (10.29)	
Pulmonary edema	536 (1.83)	124 (76)		105 (19.59)	
Cardiogenic shock	883 (3.02)	105 (56)		307 (34.77)	

Continued on next page

**Table 1 Continued**

Description	Prevalence n (%)	Door-to-Balloon Time		Mortality	
		Median, min (Quartile Range)	p	n (%)	p
First 12-lead ECG					
Anterior/septal location	10,631 (36.38)	104 (55)	<0.0001	656 (6.17)	<0.0001
No. of leads with ST-segment elevation			<0.0001		<0.0001
Left bundle branch block	580 (1.98)	132 (84)		85 (14.66)	
2	3,174 (10.86)	118 (67)		129 (4.06)	
3 and 4	19,609 (67.10)	101 (52)		728 (3.71)	
5 and more	5,565 (19.04)	97 (48)		367 (6.59)	
Unknown	294 (1.01)	109 (62)		20 (6.80)	
First 12-lead ECG: ST-segment depression	13,758 (47.08)	99 (50)	<0.0001	584 (4.24)	0.0190
Nonspecific ST or T-wave deltas	856 (2.93)	114 (71)	<0.0001	31 (3.62)	0.1867
Q-wave (acute infarct zone)	3,565 (12.20)	106 (57)	<0.0001	196 (5.50)	0.0037
Times					
Symptom onset-to-door			<0.0001		0.0073
≤1 h	9,487 (32.47)	96 (50)		462 (4.87)	
>1–2 h	8,557 (29.28)	99 (50)		339 (3.96)	
>2 h	11,178 (38.25)	110 (61)		528 (4.72)	
Time/weekday of the admission day			<0.0001		0.3824
Weekday 8:00AM–3:59PM	10,916 (37.36)	90 (52)		492 (4.51)	
Weekday 4:00PM–11:59PM	5,690 (19.47)	105 (51)		280 (4.92)	
Weekday 12:00AM–7:59AM	4,827 (16.52)	109 (56)		198 (4.10)	
Weekend 8:00AM–3:59PM	3,639 (12.45)	110 (55)		160 (4.40)	
Weekend 4:00PM–11:59PM	2,412 (8.25)	109 (49)		120 (4.98)	
Weekend 12:00AM–7:59AM	1,738 (5.95)	119 (54)		79 (4.55)	
Calendar time			<0.0001		0.0531
First year	7,978 (27.30)	105 (57)		394 (4.94)	
Second year	7,238 (24.77)	102 (54)		344 (4.75)	
Third year	7,791 (26.66)	100 (53)		339 (4.35)	
Fourth year	6,215 (21.27)	102 (52)		252 (4.05)	

ECG = electrocardiogram.

transferred out and assumed they survived to discharge. To evaluate the potential effect of decreasing length of stay on our results, we also performed secondary analyses that evaluated mortality within 72 h. The results of both of these secondary analyses were not substantially different from the original.

Statistical analyses were performed using SAS version 9.1 (SAS Institute Inc., Cary, North Carolina), HLM 5.04 for Windows (SSI, Lincolnwood, Illinois), and Stata version 8.0 (Stata Corp., College Station, Texas). The investigators had full access to all of the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

## RESULTS

**Sample characteristics.** The cohort was predominantly male (71%) and white (86%), with a mean age of 61.6 years (Table 1). A substantial proportion of patients had a prior diagnosis of coronary artery disease and/or traditional cardiac risk factors. Almost 10% received a prehospital ECG, 94% had chest pain, 11% were in overt heart failure, and 2% had left bundle branch block. A total of 62% presented within 2 h of symptom onset. There were 58% of the patients who had an ACC/AHA high-risk feature (36% with anterior/septal location, 19% with diabetes mellitus,

12% with heart rate  $\geq 100$  beats/min, and 10% with systolic blood pressure  $< 100$  mm Hg).

**Association with in-hospital mortality.** In unadjusted analysis, many patient characteristics were significantly associated with both door-to-balloon time and in-hospital mortality (Table 1). Notable exceptions were race/ethnicity and time/day of presentation. In hierarchical multivariable analysis, many patient characteristics continued to be significantly associated with mortality (Table 2). In particular, all four ACC/AHA high-risk factors were associated with increased mortality. In contrast, symptom onset-to-door time was not significantly associated with mortality.

**Door-to-balloon time and mortality.** In-hospital mortality increased significantly with increasing door-to-balloon times (Fig. 1). This relationship was seen in patients regardless of symptom onset-to-door time (Fig. 2). The association between shorter door-to-balloon times and lower mortality was seen both for patients with ACC/AHA high-risk factors and for those without these risk factors (Fig. 3). In hierarchical multivariable analysis, the odds of in-hospital mortality increased with increasing door-to-balloon time for all subgroups, whether by symptom onset-to-door time (Fig. 4A) or presence or absence of risk factors (Fig. 4B).

T1

T2

F1

F2

F3

F4


**Table 2.** Factors Independently Associated With In-Hospital Mortality in Multivariate Hierarchical Logistic Regression Model (p < 0.01)

Description	Odds Ratio (95% Confidence Interval)	p
<b>Demographics</b>		
Age		
<65 yrs	0.41 (0.36-0.47)	<0.0001
>80 yrs	1.98 (1.72-2.27)	<0.0001
Female	1.27 (1.14-1.42)	<0.0001
<b>Medical history</b>		
Current smoker	0.74 (0.64-0.84)	<0.0001
Chronic renal insufficiency	1.70 (1.36-2.13)	<0.0001
Diabetes	1.38 (1.22-1.57)	<0.0001
Hypercholesterolemia	0.63 (0.56-0.72)	<0.0001
Family history of coronary artery disease	0.82 (0.71-0.94)	0.0056
Percutaneous coronary intervention	0.74 (0.63-0.88)	0.0005
Coronary artery bypass	1.40 (1.14-1.72)	0.0014
Stroke	1.44 (1.20-1.74)	0.0001
<b>Presentation</b>		
Chest pain at presentation		
No	3.63 (3.13-4.20)	<0.0001
Unknown	3.76 (2.76-5.12)	<0.0001
Systolic blood pressure		
<100 mm Hg	3.75 (3.27-4.30)	<0.0001
>180 mm Hg	0.43 (0.33-0.55)	<0.0001
Unknown	4.59 (2.23-9.46)	<0.0001
Heart rate		
<50 beats/min	0.84 (0.69-1.03)	0.0992
>100 beats/min	2.31 (2.02-2.64)	<0.0001
Unknown	2.27 (1.08-4.75)	0.0299
First assessment of heart failure		
Rales/jugular venous distension	2.19 (1.87-2.56)	<0.0001
Pulmonary edema	3.30 (2.63-4.14)	<0.0001
Cardiogenic shock	7.25 (6.13-8.58)	<0.0001
<b>First 12-lead electrocardiogram</b>		
Anterior/septal location		
No. of leads with ST-segment elevation	1.81 (1.61-2.03)	<0.0001
Left bundle branch block		
3 and 4	2.20 (1.63-2.98)	<0.0001
5 and more	0.98 (0.81-1.18)	0.8147
Unknown	1.44 (1.18-1.77)	0.0005
Q-wave (acute infarct zone)	1.43 (0.86-2.38)	0.1659
Q-wave (acute infarct zone)	1.24 (1.06-1.45)	0.0065
<b>Times</b>		
Door-to-balloon time per 30 min	1.08 (1.05-1.11)	<0.0001

**DISCUSSION**


In this large observational study of patients with STEMI undergoing primary PCI, we found clear evidence of increased mortality with longer door-to-balloon times. This association was seen for patients regardless of symptom onset-to-door time and for patients with and without high-risk factors. These findings support the current guideline-based recommendations for rapid PCI and provide evidence that this recommendation is valid for all patients with STEMI and presentation within 6 h of the onset of symptoms.

**Time from symptom onset to reperfusion.** Shorter time from symptom onset to the administration of fibrinolytic therapy has been consistently shown to be associated with




**Figure 1.** In-hospital mortality and door-to-balloon time; p for trend < 0.001.

lower mortality for patients with STEMI (1-3,17,18). However, a meta-analysis of randomized trials found time from symptom onset to reperfusion related to mortality for fibrinolytic therapy in all patients, but for PCI only in those treated within 2 h (10). Large single-center observational studies have found similar results (4,6). In contrast, analysis of previous patients in NRM1-1 and -2 found no significant relationship between symptom onset-to-balloon time and mortality (7). Similarly, our study of NRM1-3 and -4 did not find improved survival for patients with decreased symptom onset-to-door time after adjusting for patient characteristics. In support of our findings, myocardial salvage has been found to be related to time from symptom onset to fibrinolytic therapy but independent of time from symptom onset to balloon (19). In addition to biological explanations, methodological issues may account for the poor relationship. First, the accuracy of the time of symptom onset is limited because of patient reporting error. Patients frequently are unsure of the exact time of symptom onset and usually give an estimate. Second, patients with less certainty of time of symptom onset may be more likely to get PCI than fibrinolytic therapy because of the increased risk of bleeding for fibrinolytic therapy. Finally, some of the deaths from STEMI may occur before hospital presentation. These patients would not be entered into the registry,



**Figure 2.** In-hospital mortality and door-to-balloon time in patients stratified by symptom onset-to-door time; p for trend < 0.001 for each line.



**Figure 3.** In-hospital mortality and door-to-balloon time in patients stratified by risk factor status; p for trend < 0.001 for each line. Risk factors include anterior/septal location, diabetes mellitus, heart rate >100 beats/min, systolic blood pressure <100 mm Hg.


and their absence likely dilutes the relationship between symptom onset-to-door time and mortality.

**Door-to-balloon time.** We evaluated a subset of the symptom onset-to-balloon time, the door-to-balloon time. In addition to improving the accuracy of estimate, door-to-balloon time is easier to influence because it is more


under the control of individual hospitals and physicians than symptom onset-to-door time (20). As with symptom onset-to-balloon time, prior studies evaluating the association between door-to-balloon time and mortality have had mixed results. In the Global Use of Strategies to Open Occluded Arteries in Acute Coronary Syndromes (GUSTO) IIb trial, 30-day mortality rates increased progressively with time from randomization to balloon inflation, a close surrogate for door-to-balloon time (8). Analysis of patients in a prior cohort of NRMIs also found increasing mortality with door-to-balloon times (7). Our study confirms this association in patients with more recent data as well as in various subgroups of patients.

In contrast, a recent study finding that symptom onset-to-balloon time was an independent predictor of mortality failed to find a similar relationship between door-to-balloon time and mortality (4). The discrepancy between these findings and those of our study may be explained by the fact that only 11% of patients in this single-center study had door-to-balloon times >90 min. In contrast, a majority of patients in the NRMIs registry had door-to-balloon times in excess of 90 min. A low number of patients with times >90 min may decrease sensitivity of finding a relationship. In

**A**



**B**



**Figure 4.** Independent effect of door-to-balloon time on in-hospital mortality in subgroups by (A) symptom onset-to-door time and (B) risk factor status. Reference group: door-to-balloon time <90 min. DTB = door-to-balloon; MI = myocardial infarction; SBP = systolic blood pressure.

one study, time to reperfusion was found to be important only in those at high risk (5). Our study found increasing door-to-balloon time to be related to increasing mortality for all risk groups. The magnitude of the mortality depended on the baseline risk, but the relationship with time did not differ.

**Study limitations.** Although this database is large and has been found to be reasonably generalizable (16), there are limitations. First, as mentioned previously, the time from symptom onset is obtained from the patient and may not be accurate. However, a more accurate time from symptom onset likely would not affect the main conclusions regarding door-to-balloon time. Second, there may be other risk factors that we did not examine that could identify a subgroup of patients in which door-to-balloon time is not important. Third, most of the patients were treated with door-to-balloon times greater than guideline recommendations. The importance of further reductions beyond 90 min has not been clarified. Fourth, these results cannot be extended to patients transferred from one hospital to another. Finally, door-to-balloon time may be a proxy for general quality of care, with the relationship with mortality reflecting unobserved quality measures.

**Study implications.** Efforts should continue to decrease the door-to-balloon time for all patients with STEMI undergoing primary PCI. Degree of urgency should not depend on time of symptom onset or baseline risk factors.

---

**Reprint requests and correspondence:** Dr. Harlan M. Krumholz, Yale University School of Medicine, 333 Cedar Street, PO Box 208088, New Haven, Connecticut 06520. E-mail: harlan.krumholz@yale.edu.

---

## REFERENCES

1. Fibrinolytic Therapy Trialists' (FTT) Collaborative Group. Indications for fibrinolytic therapy in suspected acute myocardial infarction: collaborative overview of early mortality and major morbidity results from all randomised trials of more than 1000 patients. *Lancet* 1994;343:311-22.
2. Newby LK, Rutsch WR, Califf RM, et al. Time from symptom onset to treatment and outcomes after thrombolytic therapy. GUSTO-1 Investigators. *J Am Coll Cardiol* 1996;27:1646-55.
3. Goldberg RJ, Mooradd M, Gurwitz JH, et al. Impact of time to treatment with tissue plasminogen activator on morbidity and mortality following acute myocardial infarction (the Second National Registry of Myocardial Infarction). *Am J Cardiol* 1998;82:259-64.
4. De Luca G, Suryapranata H, Zijlstra F, et al., ZWOLLE Myocardial Infarction Study Group. Symptom-onset-to-balloon time and mortality in patients with acute myocardial infarction treated by primary angioplasty. *J Am Coll Cardiol* 2003;42:991-7.
5. Antoniucci D, Valenti R, Migliorini A, et al. Relation of time to treatment and mortality in patients with acute myocardial infarction undergoing primary coronary angioplasty. *Am J Cardiol* 2002;89:1248-52.
6. Brodie BR, Stuckey TD, Wall TC, et al. Importance of time to reperfusion for 30-day and late survival and recovery of left ventricular function after primary angioplasty for acute myocardial infarction. *J Am Coll Cardiol* 1998;32:1312-9.
7. Cannon CP, Gibson CM, Lambrew CT, et al. Relationship of symptom-onset-to-balloon time and door-to-balloon time with mortality in patients undergoing angioplasty for acute myocardial infarction. *JAMA* 2000;283:2941-7.
8. Berger PB, Ellis SG, Holmes DR Jr., et al. Relationship between delay in performing direct coronary angioplasty and early clinical outcome in patients with acute myocardial infarction: results from the Global Use of Strategies to Open Occluded Arteries in Acute Coronary Syndromes (GUSTO-IIb) trial. *Circulation* 1999;100:14-20.
9. Brodie BR, Stone GW, Morice MC, et al. Importance of time to reperfusion on outcomes with primary coronary angioplasty for acute myocardial infarction (results from the Stent Primary Angioplasty in Myocardial Infarction Trial). *Am J Cardiol* 2001;88:1085-90.
10. Zijlstra F, Patel A, Jones M, et al. Clinical characteristics and outcome of patients with early (<2 h), intermediate (2-4 h) and late (>4 h) presentation treated by primary coronary angioplasty or thrombolytic therapy for acute myocardial infarction. *Eur Heart J* 2002;23:550-7.
11. Antman EM, Anbe DT, Armstrong PW, et al. ACC/AHA guidelines for the management of patients with ST-elevation myocardial infarction: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee to Revise the 1999 Guidelines for the Management of Patients with Acute Myocardial Infarction). *Circulation* 2004;110:e82-92.
12. Ryan TJ, Antman EM, Brooks NH, et al. 1999 update: ACC/AHA guidelines for the management of patients with acute myocardial infarction: executive summary and recommendations: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee on Management of Acute Myocardial Infarction). *Circulation* 1999;100:1016-30.
13. McNamara RL, Herrin J, Bradley EH, et al. Hospital improvement in time to reperfusion in patients with acute myocardial infarction, 1999-2002. *J Am Coll Cardiol* 2006;47:45-7.
14. French WJ. Trends in acute myocardial infarction management: use of the National Registry of Myocardial Infarction in quality improvement. *Am J Cardiol* 2000;85:5B-9B; discussion 10B-2B.
15. Rogers WJ, Bowlby LJ, Chandra NC, et al. Treatment of myocardial infarction in the United States (1990 to 1993). Observations from the National Registry of Myocardial Infarction. *Circulation* 1994;90:2103-14.
16. Every NR, Frederick PD, Robinson M, Sugarman J, Bowlby L, Barron HV. A comparison of the National Registry of Myocardial Infarction 2 with the Cooperative Cardiovascular Project. *J Am Coll Cardiol* 1999;33:1886-94.
17. Gruppo Italiano per lo Studio della Sopravvivenza nell'Infarto Miocardico. GISSI-2: a factorial randomised trial of alteplase versus streptokinase and heparin versus no heparin among 12,490 patients with acute myocardial infarction. *Lancet* 1990;336:65-71.
18. The European Myocardial Infarction Project Group. Prehospital thrombolytic therapy in patients with suspected acute myocardial infarction. *N Engl J Med* 1993;329:383-9.
19. Schomig A, Ndrepepa G, Mehilli J, et al. Therapy-dependent influence of time-to-treatment interval on myocardial salvage in patients with acute myocardial infarction treated with coronary acute stenting or thrombolysis. *Circulation* 2003;108:1084-8.
20. Luepker RV, Raczynski JM, Osganian S, et al. Effect of a community intervention on patient delay and emergency medical service use in acute coronary heart disease: the Rapid Early Action for Coronary Treatment (REACT) trial. *JAMA* 2000;284:60-7.