

CHD Clinical Practice Algorithm: PFO/Atrial Septal Defect < 5 Years of Age^{8,10,13,14}



CHD Clinical Practice Algorithm: PFO/Atrial Septal Defect 5-18 Years of Age^{8,10,13}




*Consider intervention with:


- Dilated right sided structures
- History of stroke or prothrombotic state

CHD Clinical Practice Algorithm:

Atrial Septal Defect >18 Years of Age^{11,12}



CHD Clinical Practice Algorithm: Atrial Septal Defect Post-Intervention¹⁰



- 1. ACC/AHA/ASE/HRS/ISACHD/SCAI/SCCT/SCMR/SOPE 2020 appropriate use criteria for multimodality imaging during the follow-up care of patients with congenital heart disease: a report of the American College of Cardiology Solution Set Oversight Committee and Appropriate Use Criteria Task Force, American Heart Association, American Society of Echocardiography, Heart Rhythm Society, International Society for Adult Congenital Heart Disease, Society for Cardiovascular Angiography and Interventions, Society of Cardiovascular Computed Tomography, Society for Cardiovascular Magnetic Resonance, and Society of Pediatric Echocardiography. *J Am Coll Cardiol* 2020;75:657-703.
- 2. Meijboom F, Hess J, Szatmari A, et al. Long-term follow-up (9 to 20 years) after surgical closure of atrial septal defect at a young age. *Am J Cardiol* 1993;72:1431-34.
- 3. Roos-Hesselink JW, Meijboom FJ, Spitaels SEC, at al. Excellent survival and low incidence of arrhythmias, stroke and heart failure long-term after surgical ASD closure at young age. A prospective follow-up study of 21-33 years. *Eur Heart J* 2003;24:190-97.
- 4. Cuypers JAAE, Opić P, Menting ME, et al. The unnatural history of an atrial septal defect: longitudinal 35 year follow up after surgical closure at young age. *Heart* 2013;99:1346-52.
- 5. Brassard M, Fouron JC, van Doesburg NH, Mercier LA, De Guise P. Outcome of children with atrial septal defect considered too small for surgical closure. *Am J Cardiol* 1999;83:1552-55.
- 6. Jalal Z, Hascoët S, Gronier C, et al. Long-term outcomes after percutaneous closure of ostium secundum atrial septal defect in the young: a nationwide cohort study. *JACC Cardiovasc Interv* 2018;11:795-804.
- 7. Alnasser S, Lee D, Austin PC, et al. Long term outcomes among adults post transcatheter atrial septal defect closure: systematic review and meta-analysis. *Int J Cardiol* 2018;270:126-32.
- 8. Campbell M. Natural history atrial septal defect. Br Heart J 1970;32:820-26.
- 9. Hanslik A, Pospisil U, Salzer-Muhar U, Greber-Platzer S, Male C. Predictors of spontaneous closure of isolated secundum atrial septal defect in children: a longitudinal study. *Pediatrics* 2006;118:1560-65.
- 10. McMahon CJ, Feltes TF, Fraley JK, et al. Natural history of growth of secundum atrial septal defects and implications for transcatheter closure. *Heart* 2002;87:256-69.
- 11. Baumgartner H, De Backer J, Babu-Narayan SV, et al. 2020 ESC guidelines for the management of adult congenital heart disease: the task force for the management of adult congenital heart J 2021;42:563-645.
- 12. Stout KK, Daniels CJ, Aboulhosn JA, et al. 2018 AHA/ACC guideline for the management of adults with congenital heart disease: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *J Am Coll Cardiol* 2019;73:1494-1563.
- 13. Radzik D, Davignon A, van Doesburg N, Fournier A, Marchand T, Ducharme G. Predictive factors for spontaneous closure of atrial septal defects diagnosed in the first 3 months of life. *J Am Coll Cardiol* 1993;22:851-53.
- 14. Ziebell DS, Ghaleb S, Anderson J, Statile CJ. Resource utilization in pediatric patients with secundum atrial septal defects. *Cardiol Young* 2020;30:383-87.