

Stable Ischemic Heart Disease

2016

Jane A. Linderbaum MS, CNP, AACC


Mayo Medical Center – Rochester, MN

Assistant Professor of Medicine

Associate Medical Editor AskMayoExpert

linderbaum.jane@mayo.edu

NO DISCLOSURES



Jane A. Linderbaum MS, CNP, AACCC

NO DISCLOSURES

Stable Ischemic Heart Disease Objectives

1. Identify guideline recommendations for the treatment of SIHD
2. Describe the optimal medical management for patients with SIHD
3. Define optimal therapies for special groups of patients with SIHD

Scope of the problem....

Ischemic Heart Disease



- Over 17 million people in the United States have IHD
- Up to 20% of patients with stable angina become unstable within one year
- IHD is the number one cause of death in both men and women - - over 400,000 deaths in the US annually

2012 ACC AHA Guidelines



JACC

JOURNAL of the AMERICAN COLLEGE of CARDIOLOGY

SEARCH

Advanced Search



[Home](#)

[Current Issue](#)

[All Issues](#)

[Online Before Print](#)

[Topic Collections](#)

[JACC Perspectives](#)

[Volume 60, Issue 24, December 2012 >](#)

Practice Guideline | December 2012

2012 ACCF/AHA/ACP/AATS/PCNA/SCAI/STS Guideline for the Diagnosis and Management of Patients With Stable Ischemic Heart Disease

A Report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines, and the American College of Physicians, American Association for Thoracic Surgery, Preventive Cardiovascular Nurses Association, Society for Cardiovascular Angiography and Interventions, and Society of Thoracic Surgeons

FREE

Stephan D. Fihn, MD, MPH; Julius M. Gardin, MD; Jonathan Abrams, MD; Kathleen Berra, MSN, ANP; James C. Blankenship, MD; Apostolos P. Dallas, MD; Pamela S. Douglas, MD; InAnne M. Emdin, MD; Thomas C. Gerber, MD, PhD; Alan I. Hinderliter, MD

Some tools below are only available to our subscribers or users with an online account.



PDF



Email



Share



Get Citation



Get Permissions



Article Alerts




Slideset (.ppt)



Supplemental Content



Advertisement



Internet | Protected Mode: Off

Stable

Chronic

- **Precipitating event**
- **Predictable**
- **Reproducible**
- **Unchanged**

Unstable

Rest, crescendo, ACS

- **Abrupt onset**
- **Unpredictable**
- **Unstable intensity**
- **New or changed in characteristics**

Old Cart

Onset

Location

Duration

Characteristics

Aggravating

Relieving

Treatment



Stable Angina Gone Wrong

Exacerbating Factors

- **Fever**
- **Infection**
- **Anemia**
- **Thyrotoxicosis**
- **Tachycardia**
- **Drugs that activate the Sympathetic Nervous System (↑O₂ Demand)**

CCS Angina Classification

Canadian Cardiovascular Society

- **Class I** – angina with unusually strenuous activity (>7 mets)
(no symptoms with normal activities)
- **Class II** – angina with prolonged, activity more than norm,
after meals (>5 mets)
(walking uphill, rapid stairs, emotional or cold exacerbates)
- **Class III** – angina with normal activities of daily living (2
mets)
(1-2 blocks on level, steady pace in normal conditions)
- **Class IV** – rest angina (<2 mets)
(symptoms with any activity, and sometimes rest)


Spectrum of Angina

Stable Angina

Unstable Angina

NSTEMI

STEMI



Outpatient
Management

Evaluation
and Urgent / Emergent
Treatment



Key Points – Testing / Decisions



Normal exercise test (nuclear or a normal exercise stress) is associated with a very low annual risk of cardiac death and MI

Activity 30-60 minutes of moderate-intensity aerobic activity, 5-7 days per week always and forever

Shared decision making - Patient Centered Care!

Key Points – Coronary CT



Coronary CT(CTA) is reasonable for patients with an intermediate pretest probability and:

- symptoms with prior normal test findings
- inconclusive ex/stress testing, or unable to ex/stress

Coronary CTA should not be performed for assessment of native coronary arteries with known moderate or severe calcification, or with coronary stents <3 mm in diameter in patients with known SIHD who have new or worsening symptoms not consistent with unstable angina, irrespective of ability to exercise.

Intermediate Pretest Probability

Who are they?

We need to know.....

Noninvasive Risk Stratification

High Risk (>3% annual mortality rate)

1. Severe resting LV dysfunction (LVEF <35%)
2. High-risk Duke treadmill score (score \leq -11)
3. Severe exercise-induced LV dysfunction (exercise LVEF <35%)
4. Stress-induced large perfusion defect (particularly if anterior)
5. Stress-induced multiple perfusion defects of moderate size
6. Large, fixed perfusion defect with LV dilation or increased lung uptake (thallium-201)
7. Stress-induced moderate perfusion defect with LV dilation or increased lung uptake (thallium-201)
8. Echocardiographic wall-motion abnormality (involving >2 segments) developing at a low dose of dobutamine (\leq 10 mg/kg/min) or at a low heart rate (<120 bpm)
9. Stress echocardiographic evidence of extensive ischemia

Intermediate Risk (1-3% annual mortality rate)

1. Mild/moderate resting LV dysfunction (LVEF 35-49%)
2. Intermediate-risk Duke treadmill score (-11 < score < 5)
3. Stress-induced moderate perfusion defect without LV dilation or increased lung intake (thallium-201)
4. Limited stress echocardiographic ischemia with a wall-motion abnormality only at higher doses of dobutamine involving \leq 2 segments

Low Risk (<1% annual mortality rate)

1. Low-risk Duke treadmill score (score \geq 5)
2. Normal or small myocardial perfusion defect at rest or with stress^a
3. Normal stress echocardiographic wall motion or no change of limited resting wall motion abnormalities during stress^a

SIHD 2013

Key Points - Treatment



CABG vs PCI complete revascularization is accomplished less often in patients receiving PCI than CABG, later outcomes thereafter remain unclear

Coronary artery bypass grafting (**CABG**) surgery is **probably recommended** in preference to percutaneous coronary intervention (PCI) to improve survival in some patients.

Medications SIHD

Medication - Key Points for practice



Aspirin 75-162 mg daily forever

Beta-blocker therapy started and continued for 3 years* in all patients with normal left ventricular function after MI or acute coronary syndrome

*Bangalore, Circ CV Outcomes Nov 2014

Acel – all patients with SIHD who also have hypertension, diabetes mellitus, left ventricular ejection fraction of 40% or less, or chronic kidney disease, unless contraindicated

Ranolazine can be useful when prescribed as a substitute for beta-blockers.

Ranolazine

- Metabolic modulator exact mechanism remains unknown, but lesser effect on hemodynamics
- Can be used in conjunction with ACE I, BB, CCB, nitrates in patients with inadequate symptom control
- Diabetics, intolerant to BB
- May prolong QT <http://www.crediblemeds.org/>

Moderate dose statin therapy can lower LDL-C by approximately 30-50%

- Atorvastatin 10-20 mg
- Rosuvastatin 5-10 mg
- Simvastatin 20-40 mg
- Pravastatin 40-80 mg
- Lovastatin 40 mg
- Fluvastatin XL 80 mg
- Fluvastatin 40 mg bid

Key Points



- Review medication profile
- Monitor symptoms
- Stress test only as indicated by symptoms
- Modify risk factors (continue OMT)

Patient Decision Aids

<http://shareddecisions.mayoclinic.org/>

MAYO CLINIC

Request an Appointment
Find a Doctor
Find a Job
Make a Donation

Log in to Patient Account
Translated Content

[Twitter](#) [Facebook](#) [Google+](#) [YouTube](#) [Pinterest](#)

PATIENT CARE & HEALTH INFO | DEPARTMENTS & CENTERS | RESEARCH | EDUCATION | FOR MEDICAL PROFESSIONALS | PRODUCTS & SERVICES | GIVING TO MAYO CLINIC

Shared Decisions

Home | Decision Aids | Blog | KER Unit | Patient Advocacy | Resources | Contact Us

- What are Decision Aids
- Cardiovascular Prevention DAs (Statin & Aspirin)
- Depression Medication Choice
- Diabetes Medication Decision Aid
- Osteoporosis Decision Aid
- PCI Choice

Search our site

Search entire site...

Practical Application

Optimal Medical Therapy (OMT) for SIHD



1. Lifestyle interventions
2. Pharmacologic therapy and secondary prevention
3. Treatment of ischemia and symptom control

Lifestyle/Risk Goals – What's it really mean?

Risk Factor

Smoking

Total dietary fat / sat fat

Dietary Cholesterol

Dietary sodium

Fish

BMI

Blood pressure

Lipids

Non-HDL (secondary goal)

Diabetes

Goal

Cessation

<30% cals/ <7% cals

<200 mg/day

<2000 mg/day (DASH)

3 serv/wk

BMI <25

<140/90 mmHg

Statins – mod/high (risk)

<100 mg/dl if TG ≥150

A1c <7%

72 y/o male with CAD, recent Afib

HPI 72 y/o gentleman presents with a known hx of CAD, recent TEE/DCCV for AF one month ago. He denies CP, SOB and continues to exercise regularly (swimming, walking, biking).

PMH CAD-NSTEMI six years ago, PCI LAD, GERD, DJD, BPH

72 y/o male with CAD, recent Afib

Medications

ASA 81mg, warfarin, simvastatin, metoprolol ER, omeprazole, proscar

Physical Exam


HR 66 regular – 20 – 36.5 - 126/86, 132/74

CV – JVP-nl, No bruits, heart S1,S2, no S3

Lungs clear bilat Pulses full, no peripheral edema

72 y/o male with CAD, recent Afib

Presenting ECG




Spectrum of Angina

Stable Angina

Unstable Angina

NSTEMI

STEMI



Outpatient
Management

Evaluation
and Urgent / Emergent
Treatment



Key Points



- Review medication profile
- Monitor symptoms
- Stress test only as indicated by symptoms
- Modify risk factors (continue OMT)

Stable Angina Objectives

‘Need to Know Questions’

- What are the contraindications for beta blockers?
- Which patients have improved survival with CABG vs. other treatment modalities?

Coronary angiography and revascularization should be considered in patients with the following:

- Severe or progressive angina
- Angina that not adequately relieved by medical therapy
- Congestive heart failure
- Strongly positive stress test
- Recurrent symptoms (6 to 12 months) after percutaneous coronary intervention (PCI) in a coronary distribution supplying a moderate to large amount of myocardium

i.e. symptomatic patients

Special Groups Review

- Antiplatelet therapy (ie, ASA; if allergic to ASA, then clopidogrel)
- Beta-blockade in the presence of prior myocardial infarction or symptomatic or inducible ischemia
- Angiotensin-converting enzyme inhibitors in selected patients:
 - CHF or EF less than 40%
 - Diabetes
 - Hypertension not controlled by other anti-anginal
 - May be considered in any patient with documented coronary artery disease (class IIa)
- Nitrates for symptomatic ischemia

Practice Pearls Summary



Caution with aggressive BP management - excessive reduction in diastolic pressure has not been shown to improve outcomes and has been associated with an increase in mortality (reduced coronary perfusion?)

Caution with nicotine dependence therapy - worsening of existing depression and the risk for suicide with the use of varenicline (Chantex™) FDA warning.

β -blockers caution in diabetics. The adverse event profile of β -blockers may limit their use. Avoid abrupt β -blocker withdrawal should be avoided – taper 1-3 weeks

- **Absolute contraindications:** severe bradycardia, pre-existing high degree of AVB, sick sinus syndrome, and refractory heart failure.
- **Relative contraindications** include bronchospastic disease or active peripheral arterial disease (β -blockers without vasodilating properties or selective agents at low doses may be considered).

Practical Pearls



Calcium-channel blockers should be used with caution in patients who are taking cyclosporine, carbamazepine, lithium carbonate, amiodarone, or digoxin because of potential drug/drug interactions.

Ranolazine helpful in diabetics, side effects are constipation, nausea, dizziness, and headache. The incidence of syncope is less than 1%. Ranolazine is contraindicated in combination with potent inhibitors of the CYP3A4 pathway, including ketoconazole and other azole antifungal agents, macrolide antibiotics, human immunodeficiency virus (HIV) protease inhibitors, grapefruit products or juice, and diltiazem.

Revascularization outcomes remain less favorable in women, in-hospital mortality after PCI also higher for women

Self Care Plan

Symptoms you may have

- Not feeling well.
- Shortness of breath.
- Feeling tired.
- Fast heartbeat or heart palpitations.
- Aching or pain in the chest that spreads to the neck, jaw, arms, shoulders, or back.
- Feeling fullness, pressure or tightness in your chest.
- Upset stomach or nausea, heartburn or indigestion, vomiting.

Know your medications

- Know what medications you are taking and know why you are taking them.
- Take your medications as you have been told.
- Do not stop any medication unless you talk to your health care provider.

Good health choices

- Do not smoke or use tobacco in any form.
- Avoid second-hand smoke.
- Choose food low in fat, cholesterol, sugar, and salt, also called sodium.
- Control your blood pressure.
- Go to cardiac rehabilitation regularly.
- Do regular exercise. Aim for 2½ hours each week.

- If you have diabetes, keep it under control.
- Identify stress in your life and get help to manage it.
- If you feel you are depressed, ask for help.
- Ask family and friends to support you. Stay in touch with them.
- If you drink alcohol, do it in moderation.
- Only use drugs or medications that your health care provider approves or prescribes for you.

Identify Your Zone

Green Zone Stable

You are in the green zone when you:

- Feel well.
- Have no cardiac symptoms. See above for the list of cardiac symptoms.
- Are able to do your usual daily activities.
- Make good health choices. See above for the list of good health choices.

Your plan:

- Continue to follow your treatment plan.

Yellow Zone Caution

You are in the yellow zone when you:

- Don't feel well.
- Are short of breath.
- Feel tired.
- Have a fast heartbeat, chest pain or both.
- Are not able to do your usual daily activities.
- Have other symptoms that relate to your heart condition. See above for the list of symptoms.

Your plan:

- Stop what you are doing and rest.
- Take nitroglycerin as you have been told by your health care provider.
- Wait 10 to 15 minutes before you go back to your usual activities.
- Call your care provider if your symptoms happen more often or if you are worried about your symptoms.
- Be sure you make good health choices. See above for the list of good health choices.

Red Zone Alert!

You are in the red zone when:

- You still have symptoms after you take nitroglycerin.

Your plan:

- Stop what you are doing.
- Call 9-1-1 or your local emergency room.
- Take a second dose of nitroglycerin.

Stable CAD Highlights

- **ASA** if no contraindications
 - Treat **risk factors and involve the patient!**
 - **Statins** regardless of lipid levels unless contraindicated (moderate dose)
 - **Beta blockers** as first-line anginal treatment, followed by **nitrates** and **calcium channel blockers** as indicated*
 - **Refer** high risk patients
-
- *Bangalore, Circ CV Outcomes Nov 2014

Stable Ischemic Heart Disease

Objective Summary

- 1. Identify guideline recommendations for the treatment of SIHD**
- 2. Describe the optimal medical management for patients with SIHD**
- 3. Define optimal therapies for special groups of patients with SIHD**

Thank you!

Find the guideline

Google SIHD guideline

acc.org

cardiosource.com

linderbaum.jane@mayo.edu

Resources

- <http://www.acc.org/>
- <http://www.cardiosource.com/>
- <http://www.blaufuss.org/>
- O'Rourke, R.A.; Fuster, V.; Alexander, R. W.; Roberts, R.; King III, S. B.; Prystowsky, E.N.; Nash, I. S. (2005). *11th Edition Hurst's The Heart Manual of Cardiology*. McGraw-Hill Medical Publishing Division.
- Anderson, JL; Adams, CD; Antman, EA; et al. *ACC/AHA 2007 Guidelines for the Management of Patients With Unstable Angina/Non ST-Elevation Myocardial Infarction: A Report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to Revise the 2002 Guidelines for the Management of Patients With Unstable Angina/Non ST-Elevation Myocardial Infarction)*. *Circulation*. 2007;116:e148-304.
- Bangalore, Circ CV Outcomes, November 2014.

Understand indications for further testing of patients with known CAD	Pharmacological stress with either nuclear MPI or echocardiography is recommended for risk assessment in patients with SIHD who have left bundle-branch block on ECG, regardless of ability to exercise to an adequate workload	McCully - Risk Stratification	A 66 y/o woman presents with 6 months of chest discomfort, occurring with exertion but also has had fleeting chest pain. Each episode lasts several minutes and then spontaneously resolves. She is active and able to walk two flights of stairs. The frequency and duration of pain have not changed. She has a history of DM on oral hypoglycemic agents but does not have a history of HTN, hyperlipidemia or tobacco use. Her examination is normal aside from a paradoxical split S2. Her EKG shows NSR with a LBBB. A rest echo showed an EF 66% without regional wall motion abnormalities aside from the paradoxical septal contraction	Adenosine sestamibi	EKG TMET	Exercise echocardiogram	CT angiogram	Patients with a LBBB can have a false positive result with exercise imaging test due to the dyssynchronous contraction of the septum. Pharmacological stress with either nuclear MPI or echocardiography is recommended for risk assessment in patients with SIHD who have left bundle-branch block on ECG, regardless of ability to exercise to an adequate workload.	2012 ACCF/AHA/ACP/AATS/PCNA/SCAI/STS Guideline for the Diagnosis and Management of Patients With Stable Ischemic Heart Disease: Executive Summary: A Report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines, and the American College of Physicians, American Association for Thoracic Surgery, Preventive Cardiovascular Nurses Association, Society for Cardiovascular Angiography and Interventions, and Society of Thoracic Surgeons Stephan D. Fihn, MD, MPH; et al. : J Am Coll Cardiol 2012;60:e44-164;	Pharmacological stress with either nuclear MPI or echocardiography is recommended for risk assessment in patients with SIHD who have left bundle-branch block on ECG, regardless of ability to exercise to an adequate workload . (Level of Evidence: B)
---	---	-------------------------------	--	---------------------	----------	-------------------------	--------------	--	--	--